

On strong integrality properties of the perfect matching polytope

Roland Grappe^a, Mathieu Lacroix^b and Francesco Pisanu^{b,c,*}

^aUniversité Paris Dauphine, PSL Research University, LAMSADE, CNRS UMR 7243, Paris, France

^bUniversité Sorbonne Paris Nord, LIPN, CNRS UMR 7030, Villetaneuse, 93430, France

^cUniversité Catholique de Louvain, CORE, Ottignies-Louvain-La-Neuve, 1348, Belgium

ARTICLE INFO

Keywords:

Perfect matching polytope

Box-totally dual integral polyhedron

Integer decomposition property.

ABSTRACT

This paper investigates integrality properties of perfect matching polytopes, focusing on box-total dual integrality and integer decomposition properties.

We begin by characterizing the graphs whose perfect matching polytope is a slice of the nonnegative orthant, identifying these as the solid graphs introduced by de Carvalho, Lucchesi, and Murty in “On a Conjecture of Lovász Concerning Bricks: I. The Characteristic of a Matching Covered Graph” (*Journal of Combinatorial Theory, Series B*). As a result, perfect matching polytopes of solid graphs admit a compact description. Moreover, we observe that 0, 1 polytopes that are slices of the nonnegative orthant have the integer Carathéodory property. Thus, perfect matching polytopes of solid graphs have the integer Carathéodory property. This in particular unveils a new positive case of the generalized Berge-Fulkerson conjecture.

We also establish that deciding the box-total dual integrality of a perfect matching polytope can be done in polynomial time. Additionally, we characterize the conditions under which perfect matching polytopes of two fundamental graph classes, namely near-bricks and bicritical graphs, are box-totally dual integral.

1. Introduction

This paper investigates integrality properties of perfect matching polytopes, widely studied in combinatorial optimization, with a focus on box-total dual integrality and integer decomposition properties, two fundamental notions in integer programming.

Totally dual integral and box-totally dual integral systems were introduced as a versatile framework for establishing various min-max relations in combinatorial optimization [41]. A rational linear system $Ax \leq b$ is *totally dual integral* (TDI) if the minimum in the linear duality equation

$$\max\{w^T x : Ax \leq b, x \geq \mathbf{0}\} = \min\{b^T y : A^T y \geq w, y \geq \mathbf{0}\}$$

has an integer optimal solution for every integer vector w such that the optimum is finite. If a system is TDI, then the right-hand side can be chosen integer if and only if the polyhedron described by the system is integer [24].

A stronger property for a system $Ax \leq b$ is the *box-total dual integrality* (box-TDI), which holds when $Ax \leq b, \ell \leq x \leq u$ is TDI for all rational vectors ℓ and u (with possible infinite components). Classical examples of box-TDI systems are those defined by *totally unimodular* matrices, which are the matrices whose minors are all 0, ± 1 [40]. For instance, König’s Theorem [34] and the MaxFlow-MinCut Theorem of Ford and Fulkerson [26] can be deduced from the total unimodularity of incidence matrices of bipartite graphs and of directed graphs, respectively. While every polyhedron can be described by a TDI system [29], there are polyhedra that are not described by any box-TDI system. A polyhedron is *box-TDI* if it can be described by a box-TDI system. Cook [12] proved that any TDI system describing a box-TDI polyhedron is a box-TDI system.

Box-TDI systems and polyhedra have been actively studied the past three decades. Recognizing box-TDI systems is co-NP-complete [20]; the same holds for box-TDI polyhedra [10]. Box-Mengerian matroid ports are characterized in [7]. Series-parallel graphs form a class in which several polyhedra turn out to be box-TDI: a box-TDI system describes their 2-edge-connected spanning subgraph polyhedron [8] and is generalized for k -edge-connectivity [1]; [14] provides

*Corresponding author

✉ roland.grappe@lamsade.dauphine.fr (R. Grappe); lacroix@lipn.univ-paris13.fr (M. Lacroix); francesco.pisanu@uclouvain.be (F. Pisanu)

ORCID(s): 0000-0002-7093-2175 (R. Grappe); 0000-0001-8385-3890 (M. Lacroix); 0000-0003-0799-5760 (F. Pisanu)

several other box-TDI systems; [2] proves the box-TDIness of their flow cone. Regarding box-perfect graphs, which are the perfect graphs having a box-TDI stable set polytope, new classes of box-perfect graphs are introduced in [22], and a weak box-perfect graph theorem is given in [9]. Matricial and geometrical characterizations of box-TDI polyhedra can be found in [11]. Graphs for which the matching polytope is box-TDI are characterized in [21].

A polyhedron $P = \{x : Ax \leq b\}$ has the *integer Carathéodory property* if every integer point in the k -th dilation $kP = \{x : Ax \leq kb\}$ of P can be expressed as a nonnegative integer combination of affinely independent integer points of P , for all $k \in \mathbb{Z}_{>0}$. This property was introduced in [28], where it is proved that a broad class of polyhedra, including matroid base polytopes, satisfies it. This class forms a subclass of box-TDI polyhedra. Further connections between the integer Carathéodory property and box-TDIness are discussed in [11, Section 6.3].

The integer Carathéodory property implies the *integer decomposition property*, which is to ask for each integer x in kP to be the sum of k integer points of P , for each $k \in \mathbb{Z}_{>0}$. If P has the integer decomposition property, then P is integer, and every face of P inherits the integer decomposition property [40, Section 22.10]. Originally introduced in integer programming by Baum and Trotter [3], the integer decomposition property has since been studied in fields such as algebraic geometry and combinatorial commutative algebra [30].

Several classes of polyhedra are known to have the integer decomposition property, including projections of polyhedra defined by totally unimodular matrices [42], polyhedra defined by nearly totally unimodular matrices [27], certain polyhedra defined by k -balanced matrices [47], and stable set polytopes of claw-free t -perfect graphs and h -perfect line-graphs [5]. Additional connections with Fulkerson's theory of blocking and anti-blocking polyhedra are explored in [4]. The matching polytope of a bipartite graph has the integer decomposition property, since it is described by a totally unimodular matrix. This is generalized in [45] to matchings of size $k \leq \lfloor n/2 \rfloor$ of a bipartite graph with n vertices.

In a graph, a *matching* is a subset of pairwise nonincident edges, and a *perfect matching* is a matching that covers all the vertices. The *matching polytope* of a graph is the convex hull of the incidence vectors of its matchings. Similarly, the *perfect matching polytope* of a graph is the convex hull of the incidence vectors of its perfect matchings. Since the perfect matching polytope is a face of the matching polytope and as box-TDIness is preserved under taking faces, Ding, Tan, and Zang [21]'s characterization in terms of forbidden subgraphs gives sufficient conditions for the box-TDIness of the perfect matching polytope. However, as the perfect matching polytope of a subgraph need not be a face of the perfect matching polytope of the original graph, there is no characterization of its box-TDIness in terms of forbidden subgraphs.

In this paper, we report progress on the box-TDIness and other integrality properties of perfect matching polytopes.

Contributions and outline. In Section 2, we recall the results from the literature relevant to our work.

Our contributions are threefold. First, in Section 3, we characterize the graphs for which the perfect matching polytope is the intersection of its affine hull with the nonnegative orthant: these are precisely the solid graphs. This extends a result of de Carvalho et al. [17].

Second, in Section 4, we study the box-TDIness of perfect matching polytopes for two fundamental graph classes in matching theory: near-bricks and bicritical graphs. The resulting characterization involves odd intercylicity and is based on the behavior of box-TDIness under tight cut contractions. We prove that contracting a tight cut preserves box-TDIness, while the converse fails in general. However, we show that the converse holds for 2-separation cuts.

We further prove that the perfect matching polytope is box-TDI if and only if its affine hull is box-TDI. As a consequence, deciding whether the perfect matching polytope is box-TDI can be done in polynomial time.

Finally, in Section 5, we study integer decomposition properties of perfect matching polytopes. We observe that every 0, 1 polytope that is a slice of the nonnegative orthant has the integer Carathéodory property, a result that may be of independent interest. Consequently, perfect matching polytopes of solid graphs have the integer Carathéodory property. This yields a new positive case of the generalized Berge–Fulkerson conjecture of Seymour [43].

2. Preliminaries

In this section, we give the results that we shall use throughout the paper: about box-TDI polyhedra, the perfect matching polytope, matching covered graphs and their tight cut decompositions, and the affine hull of perfect matchings. Finally, we discuss differences between the box-TDIness of the matching polytope and that of the perfect matching polytope.

In this work, all considered graphs are undirected. Without loss of generality, we assume all graphs to be connected with at least one edge, as our results extend immediately to general undirected loopless graphs. A special role is played by *odd intercylic* graphs, which are the graphs having no two vertex-disjoint odd cycles.

For a given graph $G = (V, E)$ we denote by A_G the (vertex-edge) incidence matrix of G , whose v, e entry is 1 if v is an end of e , and 0 otherwise. For $C \subseteq E$, we denote by χ^C the incidence vector of C . Throughout, $\mathbf{0}$ (resp. $\mathbf{1}$) will respectively denote a zero (resp. one) entrywise vector of appropriate size.

2.1. Box-total dual integrality, equimodularity, and integer decomposition property

A matrix is *equimodular* if it has full row rank and all maximal nonzero minors are equal up to the sign. Equimodularity is characterized in terms of total unimodularity.

Theorem 1 (Heller [31]). *A full row rank $m \times n$ matrix A is equimodular if and only if $B^{-1}A$ is totally unimodular for every nonsingular $m \times m$ submatrix B of A . Equivalently, $B^{-1}A$ is a $\{0, \pm 1\}$ -matrix for every nonsingular $m \times m$ submatrix B of A .*

It is well-known that the incidence matrix of a graph is totally unimodular if and only if the graph is bipartite [32]. The following result characterizes the class of graphs whose incidence matrix is equimodular.

Theorem 2 (Chervet et al. [10]). *The incidence matrix of a connected nonbipartite graph G is equimodular if and only if G is odd intercylic.*

Box-TDI polyhedra are characterized in terms of equimodular matrices as follows. A matrix is *face-defining* for a polyhedron P if it has full row rank and describes the affine hull of some face of P .

Theorem 3 (Chervet et al. [11]). *A polyhedron is box-TDI if and only if all its face-defining matrices are equimodular. Equivalently, each of its faces admits an equimodular face-defining matrix.*

2.2. Perfect matchings

We refer to [35] for an extensive introduction about perfect matchings. We first recall the following well-known characterization of the existence of a perfect matching, where $\mathcal{O}(G)$ denotes the family of connected components of odd cardinality of the graph G .

Theorem 4 (Tutte [46]). *A graph $G = (V, E)$ has a perfect matching if and only if $|\mathcal{O}(G \setminus S)| \leq |S|$ for all $S \subseteq V$.*

When looking at the perfect matchings of a graph, one may restrict not only to connected graphs, but even to *matching covered* graphs, which are the connected graphs in which every edge belongs to a perfect matching. By definition, a matching covered graph is *2-connected*, that is, no vertex removal disconnects the graph. The following theorem of Lovász [36] characterizes matching covered graphs. A subset S of vertices of a graph G is a *barrier* if $|\mathcal{O}(G \setminus S)| = |S|$.

Theorem 5 (Lovász [36]). *A graph having a perfect matching is matching covered if and only if each barrier is composed of pairwise nonadjacent vertices.*

Let $G = (V, E)$ be a matching covered graph. For a subset X of vertices, $E(X)$ is the set of edges of G having both ends in X , and $\delta(X)$ denotes the *cut* determined by X , that is, the set of edges having precisely one end in X . The *shores* of a cut $\delta(X)$ are X and $\bar{X} = V \setminus X$. A cut is *trivial* if one of its shores is a singleton. For $X \subseteq V$, *contracting X to a single vertex x* means replacing X by a new vertex x with $\delta(x) = \delta(X)$, and we denote the resulting graph by G/X . The two graphs G/X and $G/\bar{X}$ are referred to as the two $\delta(X)$ -*contractions* of G . A cut C of G is *tight* if $|C \cap M| = 1$ for every perfect matching M of G . A cut C of G is a *separating cut* if both of the C -contractions of G are also matching covered. Every tight cut of a matching covered graph is separating, but the converse does not hold. For example, in the prism $\bar{C}_6 := K_6 \setminus E(C_6)$, which is the complement of a cycle of length 6, the set of edges contained in no triangle forms a separating cut, which is not tight since it is also a perfect matching. Two tight cuts $\delta(X)$ and $\delta(Y)$ are *laminar* if X and Y are either disjoint or one of X and Y contains the other one. A *laminar family of tight cuts* is a family of pairwise laminar tight cuts.

A graph is *solid* if it is matching covered and all its separating cuts are tight. A matching covered graph free of nontrivial tight cuts is a *brace* if it is bipartite and a *brick* otherwise. A graph is *bicritical* if removing any couple of vertices yields a graph having a perfect matching. A graph is a brick if and only if it is 3-connected and bicritical [36]. A typical example of brick is the Petersen graph [36]; other examples of solid or nonsolid braces and bricks are given in Figure 1.




Figure 1: From left to right: the brace C_4 , which is the smallest solid bipartite graph; the brick K_4 , which is the smallest solid nonbipartite graph; the brick $\overline{C}_6$, which is the smallest nonsolid brick.

For a matching covered graph, the following holds.

Theorem 6 (Edmonds et al. [25]). *Let G be a matching covered graph, and $\delta(X)$ and $\delta(Y)$ be two laminar tight cuts of G . If $\delta(Y)$ is a cut of G/X , then $\delta(Y)$ is tight for G/X .*

Let $\mathcal{F}$ be a laminar family of nontrivial tight cuts of a matching covered graph G . Note that contracting any shore of a tight cut of $\mathcal{F}$ produces a smaller matching covered graph. By Theorem 6, the cuts from $\mathcal{F}$ that remain nontrivial in the resulting graph remain tight. Repeating this process yields a graph in which all the cuts of $\mathcal{F}$ either vanish or become trivial. We refer to the graphs obtained through this process as $\mathcal{F}$ -contractions of G . See Figure 2 illustrates the $\delta(X)$ -contractions of G when $\mathcal{F} = \{\delta(X)\}$.

When $\mathcal{F}$ is an inclusionwise maximal laminar family of nontrivial tight cuts, the resulting $\mathcal{F}$ -contractions contain no nontrivial tight cuts and hence are either bricks or braces. Thus, given such an $\mathcal{F}$, a *tight cut decomposition* of G is the collection of all bricks and braces obtained as $\mathcal{F}$ -contractions [36]. A fundamental result by Lovász [36] asserts that any tight cut decomposition of G yields the same set of bricks and braces (up to edge multiplicities).

A laminar family of nontrivial tight cuts $\mathcal{F}$ is said to satisfy the *odd cycle property* if every $\mathcal{F}$ -contraction is nonbipartite. A nontrivial tight cut $\delta(U)$ has the *odd cycle property* if the family $\mathcal{F} = \{\delta(U)\}$ has the odd cycle property. For a laminar family $\mathcal{F}$ of nontrivial tight cuts and a nontrivial tight cut $\delta(U)$ that is laminar with respect to each cut in $\mathcal{F}$, we denote by $\mathcal{F}_{G/U}$ the set of cuts in $\mathcal{F}$ that remain nontrivial tight cuts in G/U .

Let $\mathfrak{F}_G$ denote the family whose elements are maximal inclusionwise laminar families of nontrivial tight cuts having the odd cycle property. Every $\mathcal{F} \in \mathfrak{F}_G$ has the same cardinality, which is the number of bricks of G minus one [36]. When $\mathfrak{F}_G = \{\emptyset\}$ and G is nonbipartite, G is called *near-brick*. Equivalently, a near-brick is a nonbipartite matching covered graph such that no tight cut has the odd cycle property. In particular, a near-brick has a single brick, and any brick is a near-brick.

In [17], the following connection between the solidity of a graph and the one of its bricks is established.

Theorem 7 (de Carvalho et al. [17]). *A matching covered graph is solid if and only if all its bricks are.*

By de Carvalho et al. [17, Lemma 2.29], odd intercylic matching covered graphs are solid, such as bipartite matching covered graphs, odd wheels, and Möbius ladders of even order [19].

2.3. The matching polytope

Given a graph $G = (V, E)$, the matching polytope of G is denoted by $P_M(G)$. The following system of inequalities describes $P_M(G)$ and is known as *Edmonds' system* [23].

$$(1) \begin{cases} x(E(U)) \leq (|U| - 1)/2, & \text{for each } U \subseteq V \text{ with } |U| \geq 3 \text{ odd,} \\ x(\delta(u)) \leq 1, & \text{for each } u \in V, \\ x \geq \mathbf{0}. \end{cases} \quad \begin{matrix} (1a) \\ (1b) \\ (1c) \end{matrix}$$




Figure 2: A matching covered graph G , and a maximal inclusionwise laminar family of tight cuts $\mathcal{F} = \{\delta(X), \delta(Y)\}$. As $\mathcal{F} \in \mathfrak{F}_G$, G has three bricks. In particular, one can see the two $\delta(X)$ -contractions G/X and $G/\bar{X}$. While $G/\bar{X}$ is already one of the bricks of G , G/X still has two bricks that one can identify by computing $G/X/Y$ and $G/X/\bar{Y}$. All of these bricks are isomorphic to K_4 , up to parallel edges.

Cunningham and Marsh [16] proved that Edmonds' system is always TDI. In [21], the authors characterized the graphs for which Edmonds' system is box-TDI. A graph H is a *fully odd subdivision* of a graph G if H is obtained from G by subdividing each edge of G into a path of odd length (possibly the length is one), where the *length* is the number of edges in the path. Since Edmonds' system is always TDI [15], its box-TDIness is equivalent to that of the underlying polytope [12], hence we can restate their result as follows.

Theorem 8 (Ding et al. [21]). *The matching polytope of a graph G is box-TDI if and only if G has no fully odd subdivision of G_1, G_2, G_3 , and G_4 of Figure 3 as a subgraph.*




Figure 3: The graphs G_i , $i = 1, 2, 3, 4$, are the subgraphs that prevent the matching polytope from being box-TDI (up to fully odd subdivision).

2.4. The perfect matching polytope

The perfect matching polytope $P(G)$ of a graph G is described by the TDI system obtained from Edmonds' system by setting (1b) to equality. However, the following system also describes the perfect matching polytope [23], and is more convenient to investigate its box-TDIness:

$$(2) \begin{cases} x(\delta(U)) \geq 1, & \text{for each } U \subseteq V \text{ with } |U| \geq 3 \text{ odd,} \\ x(\delta(u)) = 1, & \text{for each } u \in V, \\ x \geq \mathbf{0} \end{cases} \quad \begin{matrix} (2a) \\ (2b) \\ (2c) \end{matrix}$$

Note that for bipartite graphs, inequalities (2a) are redundant, and the remaining system (2b)-(2c) is box-TDI by the total unimodularity of the incidence matrix [40, Section 19.3].

Let G be a matching covered graph and $\mathcal{F} \in \mathfrak{F}_G$. We define $M_G^{\mathcal{F}}$ as the matrix whose $|V| + |\mathcal{F}|$ rows are associated with the left-hand sides of equalities (2b) and inequalities (2a) associated with the cuts of $\mathcal{F}$. When $\mathfrak{F}_G = \{\emptyset\}$, $M_G^{\mathcal{F}}$ corresponds to the vertex-edge incidence matrix.

In their seminal works, Naddef [38], Edmonds et al. [25], and Lovász [36], proved that whenever G is nonbipartite, then $M_G^{\mathcal{F}}$ has full row rank. Moreover, the maximum number of linearly independent perfect matchings is $|E| - |V| - |\mathcal{F}| + 1$. We restate the results of Naddef [38], Edmonds et al. [25], and Lovász [36] in terms of face-defining matrices.

Theorem 9 ([25, 36, 38]). *Let G be a nonbipartite matching covered graph. Then, M_G^F is face-defining for $\text{aff}(P(G))$ for every $F \in \mathfrak{F}_G$.*

In particular, the equality associated with any nontrivial tight cut is a linear combination of equalities associated with the trivial tight cuts and the tight cuts of F .

2.5. Box-TDIness: matchings VS perfect matchings

The box-TDIness of the matching polytope implies that of the perfect matching polytope, since the latter is a face of the former. However, the converse does not hold. For instance, the perfect matching polytope of the graph G_1 in Figure 3 is box-TDI — containing only a single point — while its matching polytope is not, as shown by Theorem 8.

This phenomenon also occurs for matching covered graphs: we provide below four infinite families of graphs whose perfect matching polytope is box-TDI, but whose matching polytope is not, as they contain one of the forbidden structures of Theorem 8. Indeed, by Theorem 16 and Corollary 18, the perfect matching polytope of any fully odd subdivision of the graphs G'_1 , G'_2 , G'_3 , and G'_4 of Figure 4 is box-TDI. Their matching polytope is not box-TDI, as each of them contains one of the forbidden subgraphs of Theorem 8.




Figure 4: The graphs G'_1 , G'_2 , G'_3 , and G'_4 are matching covered graphs whose perfect matching polytope is box-TDI, yet their matching polytope is not.

3. When is the perfect matching polytope a slice of $\mathbb{R}_{\geq 0}^E$?

In [18, Theorem 2.1], de Carvalho et al. characterize the class of matching covered graphs for which the perfect matching polytope can be described by the system $x(\delta(u)) = 1$ for all $u \in V$, $x \geq \mathbf{0}$: they are the solid near-bricks and the bipartite graphs. In this section, following the line of their proof, we generalize this result and prove that the perfect matching polytope of a graph is a slice of nonnegative orthant if and only if the graph is solid¹. This yields a compact description for the perfect matching polytope of solid graphs, whereas this polytope has no compact formulation in general [39].

Theorem 10. *Let $G = (V, E)$ be a matching covered graph. Then, $P(G) = \text{aff}(P(G)) \cap \mathbb{R}_{\geq 0}^E$ if and only if G is solid.*

PROOF. We prove the result for nonbipartite graphs, as for bipartite graphs it follows instantly from the total unimodularity of the incidence matrix [32]. Let $F \in \mathfrak{F}_G$. By Theorem 9, it is sufficient to prove that $P(G) = \{x : M_G^F x = \mathbf{1}, x \geq \mathbf{0}\}$ if and only if G is solid.

($\Rightarrow$) Suppose that $G = (V, E)$ is not solid. Then, there exists a separating cut C which is not tight, and hence a perfect matching M of G such that $|M \cap C| > 1$. In [17, Lemma 2.19], de Carvalho et al. proved that a cut is separating if and only if for each edge of the cut, there exists a perfect matching that contains this edge and exactly one edge in the cut. Hence, since C is separating, for every edge $e \in M$, there exists a perfect matching M_e of G including e for which $|M_e \cap C| = 1$. Then, let $p = \frac{1}{|M|-1} ((\sum_{e \in M} \chi^{M_e}) - \chi^M)$. By construction, $p \in \{x : M_G^F x = \mathbf{1}, x \geq \mathbf{0}\}$, since $M_e, e \in E$, and M intersect each tight cut exactly once by definition. But $p^\top \chi^C < 1$, and since C is separating, its shores have odd cardinality, and hence $p \notin P(G)$.

($\Leftarrow$) Let $P = \{x : M_G^F x = \mathbf{1}, x \geq \mathbf{0}\}$, suppose that $P \neq P(G)$, and let us prove that G is not solid. Since $P(G) \subseteq P$ and every integer point of P is the characteristic vector of a perfect matching of G , P has a fractional vertex $p \notin P(G)$. Then, let C be the family of cuts associated with inequalities (2a) that are not satisfied by p . By Theorem 9, every

¹After completion of this work, we became aware that this characterization also appears in [37, Corollary 7.6(i)]. We provide a proof here for completeness.

point of P satisfies to equality the inequalities (2a) associated with tight cuts. Hence, no cut in C is tight. Let $\mathcal{M}$ denote the family of all perfect matchings of G . Let $\delta(U)$ be a cut in C such that there exists no cut $C \in C$ for which $|C \cap M| \leq |\delta(U) \cap M|$ for all M in $\mathcal{M}$ and $|C \cap M'| < |\delta(U) \cap M'|$ for some $M' \in \mathcal{M}$.

Let us prove that $\delta(U)$ is separating, that is, G/U and $G/\overline{U}$ are both matching covered. This will contradict the solidity of G . Let u be the contraction of U in G/U and suppose that G/U is not matching covered. Then, by Theorems 4 and 5, there exists a node set S such that $|\mathcal{O}((G/U) \setminus S)| > |S|$ or S is a barrier with adjacent nodes. In both cases, u belongs to S , as otherwise G is not matching covered since $|U|$ is odd. Since $\bigcup_{K \in \mathcal{O}((G/U) \setminus S)} \delta(K)$ is contained in $\delta(S)$,

we have

$$\begin{aligned} \sum_{K \in \mathcal{O}((G/U) \setminus S)} p^\top \chi^{\delta(K)} &\leq p^\top \chi^{\delta(S)} \\ &= p^\top \chi^{\delta(u) \setminus E(S)} + \sum_{s \in S \setminus \{u\}} p^\top \chi^{\delta(s) \setminus E(S)} \\ &\leq p^\top \chi^{\delta(U)} + \sum_{s \in S \setminus \{u\}} p^\top \chi^{\delta(s)} \\ &< 1 + (|S| - 1), \end{aligned}$$

where the last inequality holds since p violates the inequality (2a) associated with U but satisfies (2b). Thus, there is an odd component L of $(G/U) \setminus S$ such that $p^\top \chi^{\delta(L)} < 1$. Equalities (2b) imply that L is nontrivial so $L \in C$. Note that every odd component of $(G/U) \setminus S$ is an odd component of $G \setminus (S \setminus \{u\} \cup U)$. For every perfect matching M of G , we have

$$\begin{aligned} |\delta(U) \cap M| + |\mathcal{O}((G/U) \setminus S)| - 1 &\geq |\delta(U) \cap M| + |S| - 1 \\ &= |\delta(U) \cap M| + \sum_{s \in S \setminus \{u\}} |\delta(s) \cap M| \\ &\geq |\delta(S \setminus \{u\} \cup U) \cap M| \\ &\geq |(\bigcup_{K \in \mathcal{O}((G/U) \setminus S)} \delta(K)) \cap M| \\ &= \sum_{K \in \mathcal{O}((G/U) \setminus S)} |\delta(K) \cap M| \\ &\geq |\delta(L) \cap M| + |\mathcal{O}((G/U) \setminus S)| - 1, \end{aligned}$$

where the last inequality holds because $|\delta(K) \cap M| \geq 1$ for all $K \in \mathcal{O}((G/U) \setminus S)$. This implies that $|\delta(L) \cap M| \leq |\delta(U) \cap M|$ for every perfect matching M of G .

Let us show that this is impossible. First, if $|\mathcal{O}((G/U) \setminus S)| > |S|$, then:

$$\begin{aligned} |\mathcal{O}((G/U) \setminus S)| - 1 + |M \cap \delta(U)| &> |S| - 1 + |M \cap \delta(U)| \\ &\geq |M \cap \delta(\mathcal{O}((G/U) \setminus S))| \\ &\geq |\mathcal{O}((G/U) \setminus S)| - 1 + |M \cap \delta(L)|, \end{aligned}$$

which contradicts the choice of $\delta(U)$ in C .

Otherwise, we have $|\mathcal{O}((G/U) \setminus S)| = |S|$ and, by Theorem 5, there exists an edge $e \in S$, and since G is matching covered, there exists a perfect matching M_e of G including e . So:

$$\begin{aligned} |\mathcal{O}((G/U) \setminus S)| + |M_e \cap \delta(U)| &= |S| + |M_e \cap \delta(U)| \\ &\geq |M_e \cap \delta(S)| + 2 + |M_e \cap \delta(U)| \\ &\geq |M_e \cap \delta(\mathcal{O}((G/U) \setminus S))| + 2 \\ &\geq |\mathcal{O}((G/U) \setminus S)| + |M_e \cap \delta(L)| + 2, \end{aligned}$$

and M_e contradicts the choice of $\delta(U)$ in C .

Therefore, G/U is matching covered. Similarly, $G/\overline{U}$ is matching covered. Hence, $\delta(U)$ is separating and G is not solid. $\square$

Moreover, it turns out that the box-TDIness of a perfect matching polytope is captured by that of its affine hull.

Theorem 11. *Let G be a nonbipartite matching covered graph and $\mathcal{F} \in \mathfrak{F}_G$. Then, the following statements are equivalent:*

1. $P(G)$ is box-TDI;
2. $\text{aff}(P(G))$ is box-TDI;
3. $M_G^{\mathcal{F}}$ is equimodular;
4. $P(G) = \{x : Mx = b, x \geq \mathbf{0}\}$ with M totally unimodular.

Moreover, if $P(G)$ is box-TDI, then $P(G) = \text{aff}(P(G)) \cap \mathbb{R}_{\geq 0}^E$.

PROOF. (1. $\Rightarrow$ 2.) Every face of a box-TDI polyhedron is also box-TDI [24, (10.13)].

(2. $\Leftrightarrow$ 3.) Since an affine space has a single face, this follows from Theorems 3 and 9.

(2.&3. $\Rightarrow$ 4.) By Theorem 9, $\text{aff}(P(G)) = \{x : M_G^{\mathcal{F}}x = \mathbf{1}\}$. Since $\text{aff}(P(G))$ is integer and box-TDI, so is $\text{aff}(P(G)) \cap \mathbb{R}_{\geq 0}^E$. In particular, the latter is $P(G)$. By Theorem 1, performing a basis change in the rows of an equimodular matrix yields a totally unimodular one. Thus, let B be a basis of $M_G^{\mathcal{F}}$, $M = B^{-1}M_G^{\mathcal{F}}$, and $b = B^{-1}\mathbf{1}$. Since $M_G^{\mathcal{F}}$ is equimodular, M is totally unimodular by Theorem 1, and $P(G) = \{x : Mx = b, x \geq \mathbf{0}\}$. Note that b is integer since $\mathbf{1}$ is an integer combination of the columns of $M_G^{\mathcal{F}}$.

(4. $\Rightarrow$ 1.) When M is totally unimodular, $\{x : Mx = b, x \geq \mathbf{0}\}$ is box-TDI [32]. $\square$

By Theorem 11, when $P(G)$ is box-TDI, it is the intersection of its affine hull and the nonnegative orthant. Moreover, in Statement 4, M has full row rank and is totally unimodular, so the associated system describing $P(G)$ is a *Schröder's system* [13], that is, it is a minimal TDI system describing $P(G)$.

As a consequence of Theorem 11, the box-TDIness of perfect matching polytopes can be checked in polynomial time. First, given a matching covered graph G , one can obtain a family $\mathcal{F} \in \mathfrak{F}_G$ by applying the brick decomposition algorithm of Edmonds et al. [25] and build the matrix $M_G^{\mathcal{F}}$ in polynomial time. By Statement 3 of Theorem 11, the perfect matching polytope of G is box-TDI if and only if $M_G^{\mathcal{F}}$ is equimodular. By Theorem 1, the equimodularity of the latter can be checked as follows: find a basis B of $M_G^{\mathcal{F}}$, and then test the total unimodularity of $B^{-1}M_G^{\mathcal{F}}$. This can be done in polynomial time [44]. This polynomial case stands in contrast to the general case, where determining whether a given polytope is box-TDI is co-NP-complete [10].

Corollary 12. *Deciding whether the perfect matching polytope of a matching covered graph is box-TDI can be done in polynomial time.*

4. Near-bricks and bicritical graphs

In this section, we characterize the box-TDIness of the perfect matching polytope of near-bricks using forbidden subgraphs. We also characterize the box-TDIness of perfect matching polytopes of bicritical graphs through tight cut decomposition. As a consequence of these results, we end the section by providing two specific graphs: the first highlights that a natural necessary condition for a perfect matching polytope to be box-TDI is insufficient, and the second that a characterization of box-TDI perfect matching polytopes in terms of forbidden subgraphs is impossible.

We first prove that tight cut contractions preserve the box-TDIness of the perfect matching polytope.

Lemma 13. *Let G be a matching covered graph. If $P(G)$ is box-TDI, then so is $P(G/U)$ for each tight cut $\delta(U)$.*

PROOF. Since $\delta(U)$ is tight and G is matching covered, $P(G/U)$ is the orthogonal projection of $P(G)$ onto the coordinates indexed by $\delta(U) \cup E(\overline{U})$. Such projections preserve box-TDIness [40, Page 323], and the result follows. $\square$

4.1. The case of near-bricks

Recall that a near-brick is a nonbipartite matching covered graph in which no tight cut has the odd cycle property.

Lemma 14. *Let $G = (V, E)$ be a matching covered graph and $\delta(X)$ be a tight cut of G without the odd cycle property. If G is odd intercyclic, so is G/X .*

PROOF. Suppose that G/X contains two disjoint odd cycles C_1 and C_2 . If C_1 and C_2 are cycles of G , then G is not odd intercyclic, contradicting the assumption on G . Otherwise, one of the two cycles contains the vertex u obtained from the contraction of X . Without loss of generality, suppose $u \in V(C_1)$. Since $\delta(X)$ does not have the odd cycle property and G/X is non-bipartite, $G/\overline{X}$ is bipartite. Moreover, it is matching covered since $\delta(X)$ is tight, and hence it is 2-connected. Then, there exists an even cycle C_3 of $G/\overline{X}$ containing u . In particular, $(C_1 \setminus \delta(u)) \cup C_3$ is an odd cycle of G disjoint from C_2 , which ends the proof. $\square$

Since the brick of a near-brick is obtained by contracting tight cuts without the odd cycle property, Lemma 14 implies the following.

Corollary 15. *If a near-brick is odd intercyclic, then so is its brick.*

As a consequence, we characterize which near-bricks have a box-TDI perfect matching polytope.

Theorem 16. *The perfect matching polytope of a near-brick is box-TDI if and only if the near-brick is odd intercyclic.*

PROOF. Let $G = (V, E)$ be a near-brick.

($\Rightarrow$) We equivalently prove that if G has two vertex-disjoint odd circuits, then $P(G)$ is not box-TDI. Suppose that G contains two vertex-disjoint odd circuits. Since G is a near-brick, $\mathfrak{F}_G = \{\emptyset\}$. Thus, A_G is face-defining for $\text{aff}(P(G))$, by Theorem 9. By Theorem 2, A_G is not equimodular. Thus, $P(G)$ is not box-TDI by Theorem 3.

($\Leftarrow$) Suppose that G is odd intercyclic. By Corollary 15, its brick is odd intercyclic, hence, is a solid graph by [17, Lemma 2.29]. Thus, by Theorem 10, $P(G) = \text{aff}(P(G)) \cap \{x \geq \mathbf{0}\}$. By Theorem 2 and Theorem 9, A_G is an equimodular face-defining matrix for $\text{aff}(P(G))$. Thus, $P(G) = \{x : A_G x = \mathbf{1}\} \cap \{x : x \geq \mathbf{0}\}$ is box-TDI by Theorem 3 and the definition of box-TDIness. $\square$

4.2. The case of bicritical graphs

Let u and w be two vertices of a matching covered graph G such that $G \setminus \{u, w\}$ has precisely two connected components with vertex sets U and W of even size. Then, $\delta(X)$, with $X = U \cup \{u\}$, is a 2-separation cut with respect to u and w . Similarly, $\delta(W \cup \{u\})$ is also a 2-separation cut with respect to u and w . An illustration of 2-separation cuts is given in Figure 5. Note that every 2-separation cut is tight.




Figure 5: The two 2-separation cuts with respect to u and w are drawn in red and blue. Note that u and w may be nonadjacent.

It turns out that the converse of Lemma 13 holds for 2-separation cuts.

Theorem 17. *Let $\delta(X)$ be a 2-separation cut of a nonbipartite matching covered graph G . Then, $P(G)$ is box-TDI if and only if both $P(G/X)$ and $P(G/\overline{X})$ are.*

PROOF. In this proof, we use the following notation for a matrix A whose columns are indexed by a subset of edges of a graph: $E(A)$ denotes this set of edges.

The “only if” part comes from Lemma 13, hence let us prove the “if” part.

Let u and w be two vertices such that $G \setminus \{u, w\}$ has precisely two even connected components $G[U]$ and $G[W]$. Let $X = U \cup \{u\}$. By definition, $\delta(X)$ is a 2-separation cut with respect to u and w . Suppose that $P(G/X)$ and $P(G/\overline{X})$ are box-TDI and let us show that $P(G)$ is.

We can restrict the proof to the case where $uw \in E(G)$. Indeed, suppose the contrary. Since G is matching covered and $\delta(X)$ is tight, both G/X and $G/\overline{X}$ contain the edge uw . Adding or removing edge duplicates preserves box-TDIness, so the box-TDIness of $P(G/X)$ and $P(G/\overline{X})$ implies the one of $P((G/X) + uw)$ and $P((G/\overline{X}) + uw)$. Note that $(G/X) + uw$ and $(G/\overline{X}) + uw$ are the $\delta(X)$ -contractions of $G + uw$. Since G is matching covered so is $G + uw$. Then, proving the box-TDIness of $P(G)$ from the one of $P(G/X)$ and $P(G/\overline{X})$ can be done by proving the box-TDIness of $P(G + uw)$ from the one of $P((G/X) + uw)$ and $P((G/\overline{X}) + uw)$ as $P(G)$ is the projection onto $\mathbb{R}^{E(G)}$ of the intersection of $P(G + uw)$ with the box $\{x \in \mathbb{R}^{E(G+uw)} : x_{uw} = 0\}$. Hence, for the rest of the proof, we suppose that $uw \in E(G)$.

Consider a family $\mathcal{F} \in \mathfrak{F}_G$ such that $\mathcal{F}$ contains $\delta(X)$ if $\delta(X)$ has the odd cycle property, or such that $\mathcal{F} \cup \{\delta(X)\}$ is laminar otherwise. In both cases, the laminarity implies that the shores of the tight cuts of $\mathcal{F}$ are either contained in X or contain X . The cuts of former type correspond to rows $M_{G/\overline{X}}^{\mathcal{F}}$, the latter do not.

Let e be any edge of G . Up to replacing $U \cup \{u\}$ by $W \cup \{w\}$, we may assume that e belongs to $E(G[U \cup \{u, w\}])$. Let B be any basis of $M_G^{\mathcal{F}}$ not containing e , and denote by μ^e the e -th column of $M_G^{\mathcal{F}}$. Let us prove that there exists a $0, \pm 1$ vector x such that $Bx = \mu^e$. By Theorem 1, this will imply the equimodularity of $M_G^{\mathcal{F}}$, and hence the box-TDIness of $P(G)$ by Statement 3 of Theorem 11.

Let K and H be the graphs respectively obtained from G/X and $G/\overline{X}$ by removing edges parallel to uw . Then, e is in $E(H)$, and $P(H)$ is box-TDI as the removal of duplicate edges does not impact box-TDIness. Let B_H (resp. B_K) be the submatrix of B whose rows are indexed by $\delta(Y)$ for $Y \subseteq X$ (resp. $Y \supseteq X$) and whose columns are indexed by the edges of H (resp. K) corresponding to columns of B . Let ν^e denote the e -th column of $M_{G/\overline{X}}^{\mathcal{F}}$. Note that ν^e is the restriction of μ^e to the rows of $M_{G/\overline{X}}^{\mathcal{F}}$.

Our proof comes in two steps: we first prove that there exist a basis C_H of B_H and a vector $y \in \{0, \pm 1\}^{E(C_H)}$ such that $C_H y = \nu^e$. We then construct from y a vector $x \in \{0, \pm 1\}^{E(B)}$ such that $Bx = \mu^e$.

Step 1. First, suppose that H is bipartite. Then, $\delta(X) \notin \mathcal{F}$ and no cut of $\mathcal{F}$ has a shore contained in X . In particular, note that $[B_H, \nu^e]$ is a submatrix of A_H , and the rows of B_H are indexed by $\delta(h)$ for all $h \in V(H) \setminus \{w\}$. Since H is bipartite and connected, any matrix obtained from A_H by removing a single row, like B_H , has full row rank and is totally unimodular. Let C_H be a basis of B_H . Then, there exists a $0, \pm 1$ vector y such that $C_H y = \nu^e$.

Next, suppose that H is nonbipartite. Let M be the matrix obtained from $M_{G/\overline{X}}^{\mathcal{F}}$ by removing the duplicates of column uw . Then, B_H is the submatrix of M whose columns are indexed by those of B contained in M . Hence, a basis C_H of B_H is also a basis of M . By Theorem 9, M is face-defining for $P(H)$. Observe that M is a submatrix of $M_G^{\mathcal{F}}$. By Theorem 1 and by Statement 3 of Theorem 11, since $P(H)$ is box-TDI, there exists a $0, \pm 1$ vector y such that $C_H y = \nu^e$ for some basis C_H of B_H .

Without loss of generality, we suppose that in both cases, we choose a basis C_H such that $uw \notin E(C_H)$ if such a basis exists.

Step 2. There are two cases. First, suppose that either $uw \notin E(C_H)$ or $y_{uw} = 0$. Then, every non-null coordinate of y is associated with an edge that does not belong to G/X . Hence the associated column in B_K only contains 0. Moreover, the restriction of μ^e to the rows of B_K is the 0 vector. Defining $x \in \{0, \pm 1\}^{E(B)}$ such that $x_e = y_e$ for all $e \in E(C_H)$, and $x_e = 0$ for $e \in E(B) \setminus E(C_H)$ gives $Bx = \mu^e$.

The second case is when $uw \in E(C_H)$ and $y_{uw} \neq 0$. Then, $e \neq uw$ and every basis of B_H contains uw , from our choice of C_H and $uw \in E(C_H)$.

Consequently, there exists a basis of B_K not containing uw . Indeed, suppose by contradiction that every basis of B_K contains the edge uw . Then, removing column uw in B_K gives a matrix with rank at most $|V(K)| + |F_K| - 1$. Similarly, from our choice of C_H , since $uw \in E(C_H)$, then every basis of B_H contains uw , so removing uw from B_H gives a matrix with rank at most $|V(H)| + |F_H| - 1$. Hence, the rank of B without uw is at most $|V(G)| + |F| - 2$, which contradicts that B has full row rank and $|V(G)| + |F|$ rows.

Without loss of generality, let C_K be a basis of B_K not containing uw . Since $P(K)$ is box-TDI, there exists a $0, \pm 1$ vector z such that $C_K z = \rho^{uw}$, where ρ^{uw} is the restriction of the column of M_G^F associated with uw to the rows of B_K . Now, define the vector $\hat{x} \in \{0, \pm 1\}^E$ as follows:

$$\hat{x}_f = \begin{cases} y_f & \text{if } f \in E(C_H) \setminus \{uw\}, \\ y_{uw} z_f & \text{if } f \in E(C_K), \\ 0 & \text{otherwise.} \end{cases}$$

Note that, for $F \subseteq E(G)$, since $\hat{x}_{uw} = 0$ and since uw is in $E(C_H)$, we have:

$$\hat{x}(F) = \begin{cases} y(F \cap E(C_H)) + y_{uw} z(F \cap E(C_K)) & \text{if } uw \notin F, \\ y(F \cap E(C_H)) - y_{uw} + y_{uw} z(F \cap E(C_K)) & \text{if } uw \in F. \end{cases} \quad (3)$$

We finish the proof by showing that for each tight cut $\delta(Y)$ indexing a row of B , we have $\hat{x}(\delta(Y)) = \mu_{\delta(Y)}^e$. This immediately implies that the restriction x of $\hat{x}$ to the coordinates indexed by the columns of B satisfies $Bx = \mu^e$, since the only nonzero coordinates of $\hat{x}$ are associated with columns of $E(B)$. Let $\delta(Y)$ be a tight cut indexing a row of B .

Suppose first that $uw \notin \delta(Y)$. Then, either $Y \subseteq U$ or $Y \subseteq W$. In the former case, $\delta(Y) \cap E(C_K) = \emptyset$ so by (3), $\hat{x}(\delta(Y)) = y(\delta(Y) \cap E(C_H)) = \mu_{\delta(Y)}^e$ since $C_H y = v^e$. In the latter case, $\delta(Y) \cap E(C_H) = \emptyset$ so, by (3), $\hat{x}(\delta(Y)) = y_{uw} z(\delta(Y) \cap E(C_K)) = 0$ since $C_K z = \rho^{uw}$ and $uw \notin \delta(Y)$. Moreover, $\mu_{\delta(Y)}^e = 0$ since e is in G/X , hence not in $\delta(Y)$.

Suppose now that $uw \in \delta(Y)$. Then, we have $\rho_{\delta(Y)}^{uw} = 1$. By first applying (3), then $y(\delta(Y) \cap E(C_H)) = \mu_{\delta(Y)}^e$, and finally $y(\delta(Y) \cap E(C_K)) = \rho_{\delta(Y)}^{uw} = 1$, we obtain:

$$\hat{x}(\delta(Y)) = y(\delta(Y) \cap E(C_H)) - y_{uw} + y_{uw} z(\delta(Y) \cap E(C_K)) = \mu_{\delta(Y)}^e.$$

□

Fully odd subdividing a graph preserves the nonbox-TDIness of its matching polytope, yet the converse does not hold [21]. For perfect matchings, it is an equivalence which follows from Theorem 17.

Corollary 18. *The perfect matching polytope of a fully odd subdivision of a matching covered graph G is box-TDI if and only if $P(G)$ is.*

PROOF. Let H be the graph obtained by replacing the edge uv of G with the path $u, u_1, \dots, u_{2k}, v$, with $k \in \mathbb{Z}_{>0}$. Let $X = \{u, u_1, \dots, u_{2k}\}$. Note that $\delta(X)$ is a 2-separation cut of H . By Theorem 17, $P(H)$ is box-TDI if and only if both $P(H/X)$ and $P(H/\bar{X})$ are. Moreover, $H/X = G$ and $H/\bar{X} = C_{2k+2}$. The latter is bipartite, hence its perfect matching polytope is box-TDI. Therefore, $P(H)$ is box-TDI if and only if $P(G)$ is. □

A graph $G = (V, E)$ is *bicritical* if $G \setminus \{u, v\}$ has a perfect matching for every $u, v \in V$. Edmonds et al. [25, Section 2, page 252] — referring to $\{u, v\}$ as an articulation set — proved that a bicritical graph is a matching covered graph whose tight cut decomposition can be accomplished by a sequence of tight cut contractions stepping exclusively in 2-separation cuts. Then, Theorems 16 and 17 and Corollary 18 give the following.

Corollary 19. *The perfect matching polytope of a fully odd subdivision of a bicritical graph is box-TDI if and only if all the bricks of the graph are odd intercylic.*

An *odd wheel* is a graph composed of a circuit of odd length, and an additional vertex connected to all the vertices of the circuit. In [17], the authors proved that the only solid planar bricks are odd wheels. Thus, by Corollary 18, Theorem 7, and [17, Lemma 2.29], the following holds.

Corollary 20. *The perfect matching polytope of a fully odd subdivision of a bicritical planar graph is box-TDI if and only if all the bricks of the graph are odd wheels.*

4.3. A necessary but not sufficient condition

Lemma 13 and Theorem 16 immediately give the following.

Corollary 21. *Let G be a matching covered graph and $\mathcal{F} \in \mathfrak{F}_G$. If $P(G)$ is box-TDI, then the near-bricks obtained by any sequence contracting all the cuts of $\mathcal{F}$ are odd intercylic.*

PROOF. By Lemma 13, if $P(G)$ is box-TDI, then so is the perfect matching polytope of any near-brick that arises from a sequence of contractions of tight cuts of G . By Theorem 16, all these near-bricks are odd intercylic. $\square$




Figure 6: The Moonfish graph $\mathcal{M}$ and the unique maximal inclusionwise laminar family $\{C, C'\}$ of $\mathcal{M}$. One can check that all C -contractions and C' -contractions are odd intercylic near-bricks. Since, $\mathcal{M}$ has two bricks — K_4 and the odd wheel of order 5 — and the two C -contractions are nonbipartite, $\{C\} \in \mathfrak{F}_{\mathcal{M}}$. In particular, the two matrices above are nonsingular maximal submatrices of on the $M_{\mathcal{M}}^{(C)}$ and have —from left to right— determinant 4 and 8, up to the sign. Then, the matrix $M_{\mathcal{M}}^{(C)}$ is nonequimodular, by definition, and it is face-defining for the affine hull of $P(\mathcal{M})$ by Theorem 9. By Theorem 11, $P(\mathcal{M})$ is not box-TDI.

The converse of Corollary 21 does not hold. Figure 6 provides the Moonfish graph $\mathcal{M}$, which is a graph with two bricks illustrating this. Specifically, $\mathfrak{F}_{\mathcal{M}} = \{\{C\}, \{C'\}\}$, and all C and C' -contractions are odd intercylic. However, by Theorems 3 and 9, $P(\mathcal{M})$ is not box-TDI, as $\{C\} \in \mathfrak{F}_{\mathcal{M}}$ and $M_{\mathcal{M}}^{(C)}$ is not equimodular.

4.4. An impossible forbidden subgraph characterization

In this section, we provide a graph whose perfect matching polytope is box-TDI, and yet contains a subgraph whose perfect matching polytope is not box-TDI. This highlights that, unlike the elegant characterization of box-TDI matching polytopes given in Theorem 8, it is not possible to characterize box-TDI perfect matching polytopes solely in terms of forbidden subgraphs. This limitation arises because, in the context of perfect matchings, using a subgraph approach




Figure 7: A matching covered graph containing a fully odd subdivision of $\overline{C}_6$ (in red) as a subgraph and whose perfect matching polytope is box-TDI.

requires that if an edge is deleted, all edges that appear exclusively in perfect matchings containing the deleted edge must also be removed.

Let G be the graph illustrated in Figure 7. By Theorem 7 and [17, Lemma 2.29] G is solid, since the bricks obtained with the tight cut decomposition associated with the family

$$\mathcal{H} = \{\delta(\{1, 2, 3\}), \delta(\{5, 6, 7\}), \delta(\{8, 9, 10\}), \delta(\{11, 12, 13\}), \delta(\{14, 15, 16\})\}$$

are all K_4 . A maximal laminar subfamily of nontrivial tight cuts with the odd cycle property of $\mathcal{H}$ is

$$\mathcal{F} = \{\delta(\{1, 2, 3\}), \delta(\{5, 6, 7\}), \delta(\{8, 9, 10\}), \delta(\{11, 12, 13\})\}.$$

The family $\mathcal{F} \in \mathfrak{F}_G$ gives the matrix $M_G^{\mathcal{F}}$ which is equimodular². By Theorem 11, $P(G)$ is box-TDI.

By Theorem 16, the perfect matching polytope of the brick $\overline{C}_6$ is not box-TDI. By Corollary 18, neither are the perfect matching polytopes of its fully odd subdivisions. Since the subgraph in red in Figure 7 is a fully odd subdivision of $\overline{C}_6$, it implies that G contains a subgraph whose perfect matching polytope is not box-TDI whereas the perfect matching polytope of G is.

5. Integer decomposition properties

This section discusses consequences of the previous ones about integer decompositions properties of perfect matching polytopes, and in turn some consequences about edge-colorability.

First, we observe that $0, 1$ polytopes that are slices of the nonnegative orthant have the integer Carathéodory property. To the best of our knowledge, this statement is new and may be of independent interest. We provide a self-contained proof. Note that, in Theorem 22, $x \geq 0$ can be replaced by $x \in C$ where C is a cone.

Theorem 22. *Let $P = \{x \in \mathbb{R}^n : Ax = b, x \geq 0\}$ be a $0, 1$ polytope. Then, P has the integer Carathéodory property.*

PROOF. We prove by induction on $k \in \mathbb{Z}_{>0}$ that each integer point in kP can be expressed as a nonnegative integer sum of affinely independent integer points of P . The case $k = 1$ is immediate. Let $k \geq 2$ be integer and $x \in kP \cap \mathbb{Z}^n$, that is, x is a nonnegative integer point such that $Ax = kb$. By convexity, since $\frac{1}{k}x$ is in P , we have $x = k \sum_{w \in W} \lambda_w w$, where W is a set of vertices of P , $\sum_{w \in W} \lambda_w = 1$, and $\lambda_w > 0$ for all $w \in W$. Then, as x is a nonnegative integer and $W \subseteq \{0, 1\}^n$, we have $x \geq w$ for all $w \in W$.

Let $v \in W$ and let i be such that x_i is minimum among the coordinates x_j of x for which $v_j = 1$. Let $n_v = x_i$ and $x' = x - n_v v$. The choice of i implies $x'_i \geq 0$ and $x'_i = 0$. Note that n_v is a positive integer with $n_v \leq k$, thus x' belongs to $(k - n_v)P \cap \mathbb{Z}^n$ since $Ax = kb$ and $Av = b$.

By the induction hypothesis, $x' = \sum_{w \in W'} n_w w$ with W' composed of affinely independent integer points of P and $n_w \in \mathbb{Z}_{>0}$ for all $w \in W'$. Since $x'_i = 0$, we have $w_i = 0$ for all $w \in W'$. Therefore, $\{v\} \cup W'$ is affinely independent as $v_i = 1$. This finishes the proof since $x = n_v v + x'$. $\square$

²The equimodularity has been checked by enumerating all the nonzero maximal minors and verifying that they are equal in absolute value.




Figure 8: A nonplanar nonbipartite cubic graph with two nontrivial tight cuts T and T' , which is 3-edge-colorable by Corollary 25.

Combining Theorem 22 with Theorem 10 yields the following.

Corollary 23. *Perfect matching polytopes of solid graphs have the integer Carathéodory property.*

Therefore, perfect matching polytopes of solid graphs have the integer decomposition property. More generally, the perfect matching polytope of a matching covered graph has the integer decomposition property if and only if that of all its bricks have it, by the following remark.

Lemma 24. *Let G be a matching covered graph and $\delta(U)$ a tight cut of G . Then, $P(G)$ has the integer decomposition property if and only if both $P(G/U)$ and $P(G/\overline{U})$ have it.*

PROOF. By projection, it is immediate that contracting a tight cut preserves the integer decomposition property of the perfect matching polytope. To see the converse, let $k \in \mathbb{Z}_{\geq 0}$ and $x \in kP(G)$. Denote by $x|_{G/U}$ the restriction of x to G/U . Since $\delta(U)$ is tight, note that $x|_{G/U} \in kP(G/U)$ and $x|_{G/\overline{U}} \in kP(G/\overline{U})$. Then, if both $P(G/U)$ and $P(G/\overline{U})$ have the integer decomposition property, $x|_{G/U}$ and $x|_{G/\overline{U}}$ are respectively the sum of k perfect matchings of G/U and $G/\overline{U}$. Since the contributions of $x|_{G/U}$ and $x|_{G/\overline{U}}$ to the edges of $\delta(U) = \delta(\overline{U})$ are identical, pairing appropriately these matchings of G/U and $G/\overline{U}$ decomposes x into k perfect matchings of G . $\square$

The integer decomposition property has combinatorial consequences in terms of edge-coloring. A graph $G = (V, E)$ is d -regular if all its vertices have degree d , and it is a d -graph if in addition $|\delta(X)| \geq d$, for every $X \subseteq V$ with $|X|$ odd. Note that if G is a d -graph, deciding whether $\mathbf{1}$ belongs to the integer cone of the perfect matchings of G is equivalent to deciding whether G is d -edge-colorable. The latter means that the set of edges of G is the union of d edge-disjoint perfect matchings. A d -edge-colorable d -graph is matching covered. Moreover, observe that tight cuts in a d -edge-colorable d -graph are all of size d , and that contracting such a cut yields a d -edge-colorable d -graph. Thus, a matching covered d -graph is d -edge-colorable if and only if all its bricks are.

Lemma 24 implies that if G is a matching covered d -graph whose bricks have perfect matching polytopes with the integer decomposition property, then $P(G)$ also has the integer decomposition property, and hence G is d -edge-colorable. Deciding whether a d -regular graph is d -edge-colorable is NP-complete [33], for every integer $d \geq 3$. The generalized Berge-Fulkerson conjecture, due to Seymour [43], states that if G is a d -graph, then there exist $2d$ perfect matchings of G covering twice every edge of G . Corollary 23 implies the following positive new case of the conjecture.

Corollary 25. *If G is a solid d -graph, then G is d -edge-colorable.*

The four-color theorem is equivalent to the following: a 3-regular planar graph is 3-edge-colorable if and only if it is a 3-graph. Corollary 25 yields 3-edge-colorable 3-graphs which are non-necessarily planar graphs: for instance, Figure 8 provides a non-planar 3-edge-colorable whose bricks are K_4 's or Möbius ladders of even order.

Further questions

This paper explores integer properties of perfect matching polytopes, and Figure 9 provides an overview of the graph classes studied in this work. Several further questions are suggested by our findings.

Matching covered




Figure 9: Scheme of the graph classes studied in this paper. Bricks are shown in orange; they form a subclass of both bicritical graphs (light yellow) and near-bricks (salmon pink). Bipartite graphs, which by definition are neither near-bricks nor bicritical graphs, form another class of graphs whose perfect matching polytope is box-TDI. Solid graphs are shown in green, while graphs whose perfect matching polytope has the integer Carathéodory property (ICP) are shown in light purple. Among other examples, the figure depicts the Petersen graph, the smallest non-3-edge-colorable 3-graph, whose perfect matching polytope does not have the integer Carathéodory property.

Concerning the box-TDIness of perfect matching polytopes, the impact of barrier cuts remains to be investigated, as Edmonds et al. [25] essentially proves that every matching covered graph admits a tight cut decomposition consisting only of 2-separation cuts and barrier cuts. In this paper, we have dealt with 2-separation cuts. Thus, answering the following question would finish the characterization of the box-TDIness of the perfect matching polytope in terms of tight cut contractions: *Can one characterize which barrier cut contractions preserve the box-TDIness of perfect matching polytopes?*

Additionally, Statement 4 of Theorem 11 prompts the following question, suggesting a potential min-max theorem: *Can an explicit totally unimodular matrix be found to describe box-TDI perfect matching polytopes?*

Lastly, our findings provide a new characterization of solid graphs, though it remains unknown whether an efficient recognition algorithm or graphic characterization exists for nonplanar solid graphs [6]: *Could this new polyhedral characterization help?*

Conflict of interest

The authors declare that they have no conflicts of interest.

References

- [1] Barbato, M., Grappe, R., Lacroix, M., Lancini, E., 2023. Box-total dual integrality and edge-connectivity. *Mathematical Programming* 197, 307–336.
- [2] Barbato, M., Grappe, R., Lacroix, M., Lancini, E., Wolfier Calvo, R., 2022. The Schrijver system of the flow cone in series–parallel graphs. *Discrete Applied Mathematics* 308, 162–167.
- [3] Baum, S., Trotter, L.E., 1978. Integer rounding and polyhedral decomposition for totally unimodular systems, in: Henn, R., Korte, B., Oettli, W. (Eds.), *Optimization and Operations Research*, Springer Berlin Heidelberg, Berlin, Heidelberg. pp. 15–23.
- [4] Baum, S., Trotter, L.E., 1981. Integer rounding for polymatroid and branching optimization problems. *Siam Journal on Algebraic and Discrete Methods* 2, 416–425.
- [5] Benchetrit, Y., 2017. Integer round-up property for the chromatic number of some h-perfect graphs. *Mathematical Programming* 164, 245–262.

- [6] de Carvalho, M.H., Lucchesi, C.L., Murty, U.S.R., 2018. On tight cuts in matching covered graphs. *The Journal of Combinatorics* 9, 163–184.
- [7] Chen, X., Ding, G., Zang, W., 2008. A characterization of box-Mengerian matroid ports. *Math. Oper. Res.* 33, 497–512.
- [8] Chen, X., Ding, G., Zang, W., 2009. The box-TDI system associated with 2-edge connected spanning subgraphs. *Discrete Applied Mathematics* 157, 118–125.
- [9] Chervet, P., Grappe, R., 2024. A weak box-perfect graph theorem. *Journal of Combinatorial Theory, Series B* 169, 367–372.
- [10] Chervet, P., Grappe, R., Lacroix, M., Pisanu, F., Wolfler Calvo, R., 2023. Hard problems on box-totally dual integral polyhedra. *Discrete Optimization* 50.
- [11] Chervet, P., Grappe, R., Robert, L., 2021. Box-total dual integrality, box-integrality, and equimodular matrices. *Mathematical Programming* 188, 319–349.
- [12] Cook, W.J., 1986. On box totally dual integral polyhedra. *Mathematical Programming* 34, 48–61.
- [13] Cook, W.J., Pulleyblank, W.R., 1987. Linear systems for constrained matching problems. *Mathematics of Operations Research* 12, 97–120.
- [14] Cornaz, D., Grappe, R., Lacroix, M., 2019. Trader multiflow and box-TDI systems in series-parallel graphs. *Discrete Optimization* 31.
- [15] Cunningham, W.H., Green-Krotki, J., 1986. Dominants and submissives of matching polyhedra. *Mathematical Programming* 36, 228–237.
- [16] Cunningham, W.H., Marsh, A.B., 1978. A primal algorithm for optimum matching. *Mathematical Programming Study* 8, 50–72.
- [17] de Carvalho, M.H., Lucchesi, C.L., Murty, U., 2002. On a Conjecture of Lovász Concerning Bricks: I. The Characteristic of a Matching Covered Graph. *Journal of Combinatorial Theory, Series B* 85, 94–136.
- [18] de Carvalho, M.H., Lucchesi, C.L., Murty, U., 2004. The perfect matching polytope and solid bricks. *Journal of Combinatorial Theory, Series B* 92, 319–324.
- [19] de Carvalho, M.H., Lucchesi, C.L., Murty, U., 2006. How to build a brick. *Discrete Mathematics* 306, 2383–2410.
- [20] Ding, G., Feng, L., Zang, W., 2008. The complexity of recognizing linear systems with certain integrality properties. *Mathematical Programming* 114, 321–334.
- [21] Ding, G., Tan, L., Zang, W., 2018a. When is the matching polytope box-totally dual integral? *Mathematics of Operations Research* 43, 64–99.
- [22] Ding, G., Zang, W., Zhao, Q., 2018b. On box-perfect graphs. *Journal of Combinatorial Theory, Series B* 128, 17–46.
- [23] Edmonds, J., 1965. Maximum matching and a polyhedron with 0,1-vertices. *Journal of Research of the National Bureau of Standards* 69B.
- [24] Edmonds, J., Giles, R., 1977. A min-max relation for submodular functions on graphs. *Annals of Discrete Mathematics* 1, 185–204.
- [25] Edmonds, J., Pulleyblank, W.R., Lovász, L., 1982. Brick decompositions and the matching rank of graphs. *Combinatorica* 2, 247–274.
- [26] Ford, L.R., Fulkerson, D.R., 1956. Maximal flow through a network. *Canadian Journal of Mathematics* 8, 399–404.
- [27] Gijswijt, D., 2005. Integer decomposition for polyhedra defined by nearly totally unimodular matrices. *SIAM Journal on Discrete Mathematics* 19, 798–806.
- [28] Gijswijt, D., Regts, G., 2012. Polyhedra with the integer carathéodory property. *Journal of Combinatorial Theory, Series B* 102, 62–70.
- [29] Giles, F., Pulleyblank, W., 1979. Total dual integrality and integer polyhedra. *Linear Algebra and its Applications* 25, 191–196.
- [30] Gubeladze, J., 2023. Normal polytopes: between discrete, continuous, and random. *Journal of Pure and Applied Algebra* 227, 107187.
- [31] Heller, I., 1957. On linear systems with integral valued solutions. *Pacific Journal of Mathematics* 7, 1351–1364.
- [32] Hoffman, A.J., Kruskal, J.B., 2010. Integral boundary points of convex polyhedra, in: 50 Years of Integer Programming 1958–2008: From the Early Years to the State-of-the-Art, Springer Berlin Heidelberg, Berlin, Heidelberg. pp. 49–76.
- [33] Holyer, I., 1981. The NP-Completeness of edge-coloring. *SIAM Journal on Computing* 10, 718–720.
- [34] König, D., 1931. Gráfok és mátrixok. *Matematikai és Fizikai Lapok* 38, 116–119.
- [35] Lovász, L., Plummer, M.D., 1986. Matching theory. volume 121 of *Annals of Discrete Mathematics*, 29. North-Holland Publishing Co., Amsterdam.
- [36] Lovász, L., 1987. Matching structure and the matching lattice. *Journal of Combinatorial Theory, Series B* 43, 187–222.
- [37] Lucchesi, C.L., Murty, U.S.R., 2024. Perfect Matchings: A Theory of Matching Covered Graphs. Springer.
- [38] Naddef, D., 1982. Rank of maximum matchings in a graph. *Mathematical Programming* 22, 52–70.
- [39] Rothvoss, T., 2017. The matching polytope has exponential extension complexity. *Journal of the ACM (JACM)* 64, 1 – 19.
- [40] Schrijver, A., 1986. Theory of linear and integer programming. Wiley-Interscience series in discrete mathematics and optimization.
- [41] Schrijver, A., 2003. Combinatorial Optimization: Polyhedra and Efficiency. volume B. Springer Berlin.
- [42] Sebő, A., 2009. Path partitions, cycle covers and integer decomposition, in: Lipshteyn, M., Levit, V.E., McConnell, R.M. (Eds.), Graph Theory, Computational Intelligence and Thought: Essays Dedicated to Martin Charles Golumbic on the Occasion of His 60th Birthday. Springer Berlin Heidelberg, Berlin, Heidelberg. pp. 183–199.
- [43] Seymour, P., 1979. On multi-colourings of cubic graphs, and conjectures of fulkerson and tutte. *Proceedings of The London Mathematical Society*, 423–460.
- [44] Seymour, P.D., 1980. Decomposition of regular matroids. *Journal of Combinatorial Theory. Series B* 28, 305–359.
- [45] Torres, J.C., 2023. Normality of k -matching polytopes of bipartite graphs. *arXiv:2306.11910*.
- [46] Tutte, W.T., 1947. The factorization of linear graphs. *Journal of The London Mathematical Society-second Series*, 107–111.
- [47] Zambelli, G., 2007. Colorings of k -balanced matrices and integer decomposition property of related polyhedra. *Operations Research Letters* 35, 353–356.